

# Rapid Communication

## Cytokine-activated Human Endothelial Cells Synthesize and Secrete a Monocyte Chemoattractant, MCP-1/JE

Barrett J. Rollins,\* Teizo Yoshimura,† Edward J. Leonard,† and Jordan S. Pober‡

From the Division of Medicine,\* Dana-Farber Cancer Institute and Harvard Medical School, Boston, Massachusetts; Immunopathology Section,† Laboratory of Immunobiology, National Cancer Institute, Frederick, Maryland; and the Department of Pathology,‡ Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts

**We have demonstrated inducible expression of the mRNA encoding the monocyte chemoattractant MCP-1, the human homolog of the JE gene, in endothelial cells within 3 hours of treatment with IL-1 $\beta$  and tumor necrosis factor. IFN- $\gamma$  also induced expression of this mRNA after 24 hours, but to a lesser extent. MCP-1/JE protein steadily accumulated in the medium of endothelial cells during a 48-hour exposure to IL-1 $\beta$ . Medium conditioned by IL-1 $\beta$ -treated endothelial cells contained monocyte chemoattractant activity that was immunoadsorbed by anti-MCP-1 antibodies. These results suggest that endothelial cells secrete a monocyte chemoattractant, MCP-1/JE, in response to inflammatory mediators, and thus may contribute to the accumulation of monocytes at sites of inflammation. (Am J Pathol 136:1229–1233)**

Human vascular endothelial cells participate in acute inflammatory and immune reactions and are believed to play a role in chronic vascular abnormalities, such as atherosclerosis.<sup>1,2</sup> Local infiltration by various leukocyte populations is a prominent characteristic of these conditions. Activated vascular endothelium can contribute to inflammation by stimulating leukocyte migration. For example, cultured endothelial cells treated with IL-1<sup>1</sup> or TNF secrete the neutrophil chemoattractant NAP-1 (also called IL-8, MDNCF, NAF, or 3-10C).<sup>3</sup> A structurally related

monocyte chemoattractant, called MCP-1,<sup>4,5</sup> MCAF,<sup>6,7</sup> or SMC-CF,<sup>8,9</sup> has also been described. It has been identified in culture fluids of many mammalian cell types, including tumor cells,<sup>4</sup> smooth muscle cells,<sup>9</sup> and phytohemagglutinin-stimulated blood mononuclear cells.<sup>10</sup> The gene encoding this protein is the human homolog of the platelet-derived growth factor (PDGF)<sup>1</sup>-inducible murine gene JE.<sup>11</sup> Recently it has been shown that IL-1 $\beta$  and TNF also induce the expression of MCP-1/JE mRNA in endothelial cells,<sup>12</sup> and that minimally modified LDL induces the secretion of SMC-CF.<sup>13</sup> We show here that human endothelial cells express MCP-1/JE mRNA after treatment with IL-1 $\beta$ , TNF, and IFN- $\gamma$ , that endothelial cells secrete an immunoreactive form of MCP-1/JE protein, and that this protein accounts for most of the monocyte chemoattractant activity produced by endothelial cells in response to IL-1 $\beta$ .

### Materials and Methods

#### Cell Culture and Cytokines

Human endothelial cells were isolated from 3 to 5 umbilical veins by collagenase digestion, pooled, and serially subcultured. Cells were maintained in medium containing fetal calf serum and endothelial cell growth factor. Culture conditions and reagents have been described elsewhere.<sup>14</sup> Cytokines used in these studies included recombinant human IL-1 $\beta$  (expressed in *Escherichia coli*, 1.0  $\times$  10<sup>7</sup> U/mg, a gift of Dr. A. Shaw, Glaxo, Geneva, Switzerland), recombinant human tumor necrosis factor (TNF)

---

Supported by Public Health Service grant CA48675, awarded to Barrett J. Rollins, SO7RR05526-27 awarded to the Dana-Farber Cancer Institute, and HL36003 awarded to Jordan S. Pober, who is an Established Investigator of the American Heart Association.

Accepted for publication April 10, 1990.

Address reprint requests to Barrett J. Rollins, Dana-Farber Cancer Institute, 44 Binney St., Boston, MA 02115.

(expressed in *E. coli*,  $2.5 \times 10^7$  U/mg, a gift of Biogen, Cambridge, MA), and recombinant IFN- $\gamma$  (expressed in Chinese hamster ovary cells,  $2.5 \times 10^6$  U/ml conditioned medium, a gift of Professor W. Fiers, State University of Ghent, Belgium). All of the cytokine preparations were active in appropriate bioassays and were completely neutralized by relevant specific antisera. All experiments were conducted at subcultures 3 to 5, by which time mononuclear phagocytes were not detectable in these cultures.

### RNA Analysis

RNA was isolated from  $3$  to  $5 \times 10^6$  cells by guanidinium isothiocyanate extraction followed by centrifugation through cesium chloride.<sup>15</sup> RNA was electrophoretically fractionated through a 1.5% agarose 12.2 mol/l (molar) formaldehyde gel and transferred to nitrocellulose filters. Baked filters were hybridized at 42°C as described.<sup>16</sup> Probes were nick translated to a specific activity of more than  $10^8$  CPM/ $\mu$ g, and included *JE*, the *Xho*I fragment from *phJE-34*<sup>11</sup>;  $\gamma$ IP-10, the *Pst*I fragment from *pIFN $\gamma$ -31*,<sup>17</sup> a gift of Dr. A. Luster (Massachusetts General Hospital, Boston, MA); and NAP-1, a *Pst*I-*Eco*RI fragment from the IL-8 cDNA cloned into *pKK233-2*, a gift of Dr. S. L. Kunkel (University of Michigan Medical School, Ann Arbor, MI). Densitometry measurements were performed using an LKB Ultrascan enhanced laser densitometer (Pharmacia LKB Biotechnology Inc., Piscataway, NJ).

### Immune Precipitation

Two T75 flasks (Corning, Corning, NY) each containing  $2 \times 10^6$  confluent endothelial cells were treated with 250  $\mu$ Ci/ml [<sup>35</sup>S]-methionine (DuPont NEN, Boston, MA) in complete medium for 30 minutes, after which 10 U/ml IL-1 $\beta$  was added to one flask. At the indicated times, 0.5 ml medium was withdrawn from each flask and made 2 mmol/l (millimolar) in PMSF. Cells and debris were removed by centrifugation, an equal volume of RIPA buffer (50 mmol/l TRIS-HCl [pH 7.5], 150 mmol/l NaCl, 0.1% SDS, 1% Triton X-100, 0.5% deoxycholate, 1 mmol/l PMSF) was added, and immune precipitation carried out with anti-*JE* antiserum or control serum as described.<sup>11</sup>

### Preparation of Endothelial Cell Conditioned Medium

A total of  $50 \times 10^6$  endothelial cells were treated for 24 hours with 10 U/ml IL-1 $\beta$ . After ultrafiltration of a portion of the medium using a YM-5 filter (Amicon, Danvers, MA), it was fractionated by HPLC gel filtration using a  $7.5 \times 600$

mm TSK-2000 column (Toyo Soda, Tokyo, Japan) as described.<sup>18</sup>

### Immunoabsorption of Monocyte Chemotactic Activity

Twenty-five microliters (packed volume) of Protein A-Sepharose was rinsed with Gey's balanced salt solution (GBSS) with 0.2% BSA. Fifty micrograms anti-MCP-1 IgG<sup>18</sup> was added to the column, which was then rinsed with GBSS/0.2% BSA. Thirty-five microliters of conditioned medium was applied to the column and the unbound fraction was eluted with 70  $\mu$ l GBSS/0.2% BSA, yielding a total of 105  $\mu$ l unbound material. Control columns included normal rabbit IgG Protein A-Sepharose and Protein A-Sepharose alone.<sup>19</sup> Monocyte chemotaxis dose-response curves were generated for each of the effluent preparations; the concentration of monocyte chemotactic activity in each was defined as the reciprocal of the dilution showing half-maximal activity.<sup>4,18</sup>

### Monocyte Chemotaxis


Chemotaxis assays were performed in a 48-well microchamber apparatus, as described, using freshly isolated human peripheral blood mononuclear cells.<sup>20</sup>

## Results

### MCP-1/*JE* mRNA Expression in Endothelial Cells

Figure 1 shows that endothelial cells expressed detectable amounts of MCP-1/*JE* mRNA 3 hours after feeding them fresh medium, and that this expression decreased to nearly undetectable levels after 24 hours. After 3 hours of treatment, IL-1 $\beta$  and TNF induced sixfold greater increases in MCP-1/*JE* mRNA. The relative induction of MCP-1/*JE* by these cytokines was even greater after 24 hours of treatment (10- to 15-fold), although the absolute levels of expression appeared to be no higher than they were at 3 hours. IFN- $\gamma$  inconsistently induced MCP-1/*JE* mRNA expression at 3 hours, but as shown in Figure 1, reproducibly induced small increases at 24 hours.

MCP-1/*JE* is closely related to a family of proteins that are involved in the inflammatory response,<sup>21</sup> including in humans the IFN- $\gamma$ -inducible gene  $\gamma$ IP-10,<sup>17</sup> and the neutrophil-specific chemoattractant NAP-1.<sup>22-25</sup> Because the genes encoding these proteins are also expressed in cytokine-activated endothelial cells, we asked whether the regulation of MCP-1/*JE* expression was similar to that of



**Figure 1.** Human endothelial cells were treated as indicated and their RNA was isolated and blotted onto nitrocellulose, as described in Materials and Methods. For this figure, a single blot was examined using each indicated probe, stripping the blot between experiments. (Residual MCP-1/JE probe can be seen in the  $\gamma$ IP-10 experiment at a molecular size smaller than that of  $\gamma$ IP-10 mRNA.) Three independent experiments demonstrated similar results. NA, no addition.

$\gamma$ IP-10 or NAP-1. Figure 1 shows that of the cytokines tested, only IFN- $\gamma$  induced the expression of  $\gamma$ IP-10, while IL-1 $\beta$  and TNF induced NAP-1 expression. Unlike MCP-1/JE, however, NAP-1 expression was not significantly induced after 24 hours of treatment with IFN- $\gamma$ .


### Secretion of MCP-1/JE Protein by Endothelial Cells


As predicted by the results shown in Figure 1, cells incubated in complete medium secreted detectable amounts of MCP-1/JE protein that steadily accumulated over 48 hours (Figure 2). However, in the presence of IL-1 $\beta$ , cells

secreted MCP-1/JE earlier (detectable at 2 hours) and to higher levels than untreated cells. The maximal difference in MCP-1/JE secretion between treated and untreated cells occurred between 10 and 24 hours after IL-1 $\beta$  treatment. Although the antiserum precipitates a large number of proteins, MCP-1/JE is the only protein specifically precipitated. We have further documented the presence of MCP-1/JE protein in IL-1 $\beta$ -treated endothelial cell conditioned medium by Western blotting. In addition, we have confirmed the specificity of this antiserum by blocking its recognition of MCP-1/JE in Western blots using excess purified recombinant MCP-1/JE (data not shown).

MCP-1/JE exists in two forms, MCP-1/JE- $\alpha$  of  $M_r$  15,000, and MCP-1/JE- $\beta$  of  $M_r$  11,000,<sup>4,11</sup> recognized

**Figure 2.** Human endothelial cells were radiolabeled with [<sup>35</sup>S]methionine in the absence (NA) or presence of 10 U/ml IL-1 $\beta$ . At the indicated times, an aliquot of medium was withdrawn and subjected to immune precipitation using anti-JE antiserum or control serum (PI). Precipitated material was analyzed by electrophoresis through an SDS-containing 17.5% polyacrylamide gel. A second experiment demonstrated similar results.





**Figure 3.** A: Medium conditioned by IL-1 $\beta$ -treated human endothelial cells was concentrated and injected into a TSK-2000 HPLC gel filtration column equilibrated in PBS, and 1 ml fractions were collected at a flow rate of 1 ml/minute. Fraction aliquots were diluted in GBSS/0.2% BSA and assayed for monocyte chemoattractant activity. The column was calibrated with BSA, OVA (ovalbumin), CTA (chymotrypsinogen A), and CTC (cytochrome C). B: Medium conditioned by IL-1 $\beta$ -treated human endothelial cells was passed over columns containing rabbit anti-MCP-1 IgG, normal rabbit IgG, or protein A alone, and the effluents were tested for monocyte chemotactic activity as described in Materials and Methods. MCA, monocyte chemotactic activity. Values for MCA were 5.2 U/ml in the effluent from the rabbit IgG column, 4.2 U/ml in the protein A effluent, and 0.6 U/ml in the anti-MCP-1 effluent.

equally well by our antiserum.<sup>11</sup> In contrast to all other human cell types examined so far, which radiolabel the two forms to equal specific activities and secrete the two forms in nearly equimolar amounts, Figure 2 shows that endothelial cells predominantly secrete the  $\alpha$  form.

### Chemotactic Activity in Endothelial Cell Conditioned Medium

Medium conditioned by IL-1 $\beta$ -treated endothelial cells was chemotactic for human monocytes with maximal activity at a dilution of 1:25 (data not shown). This medium was concentrated sixfold and fractionated by HPLC gel filtration. Figure 3A shows that the monocyte chemotactic

activity was present as a single peak that eluted one fraction earlier than MCP-1/JE derived from glioma cells or fibroblasts.<sup>18</sup> To identify the chemotactic activity, fractions 18 to 20 were pooled and aliquots were applied to an anti-MCP-1 Protein A-Sepharose column and to control columns. Figure 3B shows that chemotactic activity was present in effluents from the control columns, but 88% of the chemotactic activity was removed by the anti-MCP-1 column, indicating that nearly all the monocyte chemotactic activity in the endothelial cell conditioned medium could be accounted for by MCP-1/JE.

### Discussion

We have demonstrated that human endothelial cells express little MCP-1/JE mRNA under standard culture conditions and that a low level of expression can be induced by feeding with fresh medium. Substantially higher levels can be induced by treating cells with IL-1 $\beta$  and TNF, and to a lesser extent by IFN- $\gamma$ . These results confirm and extend the preliminary observations of Strieter et al.,<sup>12</sup> and Takehara et al.<sup>26</sup> We report a profile of cytokine regulation that distinguishes MCP-1/JE from both NAP-1 and  $\gamma$ IP-10. We also demonstrate that endothelial cells secrete predominantly the  $\alpha$  form of MCP-1/JE and that this protein is the factor responsible for monocyte chemotactic activity in the medium of IL-1 $\beta$ -treated endothelial cells.

The secretion of a monocyte-specific chemoattractant by cytokine-activated endothelial cells provides part of the basis for the accumulation of monocytes at sites of vascular injury and inflammation. In addition to inducing chemoattraction by means of MCP-1/JE, IL-1, and TNF also cause increased adhesion of monocytes to the endothelial cell surface, probably by inducing the expression of specific endothelial-monocyte adhesion molecules.<sup>27</sup> A similar mechanism for the accumulation of neutrophils was provided by the demonstration of NAP-1 secretion by IL-1 $\beta$  and TNF-treated endothelial cells,<sup>3</sup> concomitant with expression of ELAM-1.<sup>28</sup> Despite the similarities between MCP-1/JE and NAP-1, there are differences in the controls governing their inducible expression in endothelial cells, such as the induction of MCP-1/JE by IFN- $\gamma$  versus the lack of induction of NAP-1 by this agent. Such differences in cytokine secretion as well as adhesion molecule expression by endothelial cells may underlie the temporally specific appearance of various leukocytes at sites of inflammation *in vivo*.

### References

1. Pober JS: Cytokine-mediated activation of vascular endothelium: Physiology and pathology. *Am J Pathol* 1988, 133: 426-433

2. Cotran RS, Pober JS: Effects of cytokines on vascular endothelium. *Kidney Int* 1989, 35:969-975
3. Strieter RM, Kunkel SL, Showell HJ, Remick DG, Phan SH, Ward PA, Marks RM: Endothelial cell gene expression of a neutrophil chemotactic factor by TNF- $\alpha$ , LPS, and IL-1 $\beta$ . *Science* 1989, 243:1467-1469
4. Yoshimura T, Robinson EA, Tanaka S, Appella E, Kuratsu JI, Leonard EJ: Purification and amino acid analysis of two human glioma-derived monocyte chemoattractants. *J Exp Med* 1989, 169:1449-1459
5. Yoshimura T, Yuhki N, Moore SK, Appella E, Lerman MI, Leonard EJ: Human monocyte chemoattractant protein (MCP-1): Full-length cDNA cloning, expression in mitogen-stimulated blood mononuclear leukocytes, and sequence similarity to mouse competence gene JE. *FEBS Lett* 1989, 244:487-493
6. Matsushima K, Larsen CG, DuBois GC, Oppenheim JJ: Purification and characterization of a novel monocyte chemotactic and activating factor produced by a human myelomonocytic cell line. *J Exp Med* 1989, 169:1485-1490
7. Furutani Y, Nomura H, Notake M, Oyamada Y, Fukui T, Yamada M, Larsen CG, Oppenheim JJ, Matsushima K: Cloning and sequencing of the cDNA for human monocyte chemotactic and activating factor (MCAF). *Biochem Biophys Res Commun* 1989, 159:249-255
8. Graves DT, Jiang YL, Williamson MJ, Valente AJ: Identification of monocyte chemotactic activity produced by malignant cells. *Science* 1989, 245:1490-1493
9. Valente AJ, Graves DT, Vialle-Valentin CE, Delgado R, Schwartz CJ: Purification of a monocyte chemotactic factor secreted by nonhuman primate vascular cells in culture. *Biochem* 1988, 27:4162-4168
10. Yoshimura T, Robinson EA, Tanaka S, Appella E, Leonard EJ: Purification and amino acid analysis of two human monocyte chemoattractants produced by phytohemagglutinin-stimulated human blood mononuclear leukocytes. *J Immunol* 1989, 142:1956-62
11. Rollins BJ, Stier P, Ernst TE, Wong GG: The human homologue of the JE gene encodes a monocyte secretory protein. *Mol Cell Biol* 1989, 9:4687-4695
12. Strieter RM, Wiggins R, Phan SH, Wharram BL, Showell HJ, Remick DG, Chensue SW, Kunkel SL: Monocyte chemotactic protein gene expression by cytokine-treated human fibroblasts and endothelial cells. *Biochem Biophys Res Commun* 1989, 162:694-700
13. Berliner JA, Valente AJ, Territo MC, Navab M, Gerrity R, Schwartz CJ, Fogelman AM: Monocyte chemotactic factors produced by endothelial cells are immunologically related to the factor produced by smooth muscle cells. *Arteriosclerosis* 1989, 9:697(Abstr)
14. Kurt-Jones EA, Fiers W, Pober JS: Membrane IL-1 induction on human endothelial cells and dermal fibroblasts. *J Immunol* 1987, 139:2317-2324
15. Chirgwin J, Przybyla A, MacDonald R, Rutter W: Isolation of biologically active ribonucleic acid from sources enriched in ribonuclease. *Biochemistry* 1979, 18:5294-5299
16. Rollins BJ, Morrison ED, Stiles CD: Cloning and expression of JE, a gene inducible by platelet-derived growth factor and whose product has cytokine-like properties. *Proc Natl Acad Sci USA* 1988, 85:3738-3742
17. Luster AD, Unkeless JC, Ravetch JV:  $\gamma$ -Interferon transcriptionally regulates an early-response gene containing homology to platelet proteins. *Nature* 1985, 315:672-676
18. Yoshimura T, Leonard EJ: Secretion by human fibroblasts of monocyte chemoattractant protein-1 (MCP-1), the product of gene JE. *J Immunol* 1990, 144:2377-2383
19. Leonard EJ, Skeel A: Disposable microliter immunoabsorbent columns: construction and operation. *J Immunol Methods* 1985, 82:341-348
20. Falk W, Goodwin RHJ, Leonard EJ: A 48-well microchemotaxis assembly for rapid and accurate measurement of leukocyte migration. *J Immunol Methods* 1980, 33:239-247
21. Wolpe SD, Cerami A: Macrophage inflammatory proteins 1 and 2: Members of a novel superfamily of cytokines. *FASEB J* 1989, 3:2563-2573
22. Yoshimura T, Matsushima K, Tanaka S, Robinson EA, Appella E, Oppenheim JJ, Leonard EJ: Purification of a human monocyte-derived neutrophil chemotactic factor that has peptide sequence similarity to other host defense cytokines. *Proc Natl Acad Sci USA* 1987, 84:9233-9237
23. Walz A, Peveri P, Aschauer H, Baggiolini M: Purification and amino acid sequencing of NAF, a novel neutrophil activating factor produced by monocytes. *Biochem Biophys Res Commun* 1987, 149:755-761
24. Schmid J, Weissman C: Induction of mRNA for a serine protease and a  $\beta$ -thromboglobulin-like protein in mitogen-stimulated human leukocytes. *J Immunol* 1987, 139:250-256
25. Matsushima K, Morishita K, Yoshimura T, Lavu S, Kobayashi Y, Lew W, Appella E, Kung HF, Leonard EJ, Oppenheim JJ: Molecular cloning of a human monocyte-derived neutrophil chemotactic factor (MDNCF) and the induction of MDNCF mRNA by interleukin 1 and tumor necrosis factor. *J Exp Med* 1988, 167:1883-1893
26. Takehara K, LeRoy EC, Grotendorst G: TGF- $\beta$  inhibition of endothelial cell proliferation: Alteration of EGF binding and EGF-induced growth-regulatory (competence) gene expression. *Cell* 1987, 49:415-422
27. Bevilacqua MP, Pober JS, Wheeler ME, Cotran RS, Gimbrone MAJ: Interleukin 1 acts on cultured human vascular endothelial cells to increase the adhesion of polymorphonuclear leukocytes, monocytes, and related leukocyte cell lines. *J Clin Invest* 1985, 76:2003-2011
28. Bevilacqua MP, Pober JS, Mendrick DL, Cotran RS, Gimbrone MAJ: Identification of an inducible endothelial leukocyte adhesion molecule, ELAM 1. *Proc Natl Acad Sci USA* 1987, 84:9238-9242

## Acknowledgments

The authors thank Teri Bakewell for cell culture assistance and Peter Stier for technical assistance.